

BYC 744 Proteomics-Mass Spectrometry

Qualitative and quantitative burrowing of the proteome

Stephen Barnes, PhD
sbarnes@uab.edu


Steve Barnes BMG 744 2-8-08

Topics

- **Qualitative changes in the predicted proteome**
 - Digging deeper - SELDI/MALDI
 - Posttranslational modifications
 - Isolation and characterization
 - Making use of the chemistry of the modification
- **Quantitative aspects**
 - Isotope labeling
 - ICAT, $^{18}\text{O}/^{16}\text{O}$, $^{15}\text{N}/^{14}\text{N}$
 - Chemical labeling
 - Absolute measures

Steve Barnes BMG 744 2-8-08


Surface enhanced laser desorption ionization (SELDI)



SELDI chip (plate) with local chemistry on surface causing selective binding of peptides


Steve Barnes BMG 744 2-8-08

MALDI-TOF analysis of peptides recovered by C₁₈ extraction of sera from prostate patients, with and without metastases, and controls



Steve Barnes BMG 744 2-8-08 [Lam et al., Proteomics 5, 2927](#)


Identification of a significantly altered peptide in men with metastatic prostate cancer



But what is it?

Steve Barnes BMG 744 2-8-08 Lam et al., Proteomics 5, 2927

Purification of prostate peptide by SPE extraction and strong cation exchange



Serum was first extracted with a C₁₈ SPE cartridge


SPE fraction was passed over a [H⁺]-form cation exchange resin. Bound peptides were eluted with increasing step gradients of NaCl

Analyses of fractions were performed by MALDI-TOF MS

The most enriched fraction was contaminated with albumin


Steve Barnes BMG 744 2-8-08 Lam et al., Proteomics 5, 2927

Purification of prostate peptide needed 2D-electrophoresis




Steve Barnes BMG 744 2-8-08 [Lam et al., Proteomics 5, 2927](#)

Peptide mass fingerprint of purified prostate cancer peptide




Steve Barnes BMG 744 2-8-08 [Lam et al., Proteomics 5, 2927](#)

Non-mass spec verification of prostate peptide



Western blot analysis for PF4



ELISA analysis for PF4

Moral: proteomics is a serious business that requires multiple dimensions of separation - glib methods don't work

Steve Barnes BMG 744 2-8-08 [Lam et al., Proteomics 5, 2927](#)

General classes of modification

- Biochemical events involving peptide processing
- Biochemical events stimulated by enzymes
- Chemical events driven by reactive species
- Chemical events determined by investigator

Steve Barnes BMG 744 2-8-08

Examples of peptide processing of polypeptides

- Head groups of membrane proteins
- The family of proteins in HIV that are manufactured as one protein and then hydrolyzed by a protease
- Chromogranin - a brain protein consisting of several bioactive peptides
- Formation of β -amyloid

Steve Barnes BMG 744 2-8-08




Figure 3. Peptide-Encoding Regions and Putative Functional Domains of Human Chromogranin A (CgA).

Arabic numbers designate amino acids in the mature protein (minus signal peptide). Roman numerals designate exon numbers. The intron-exon structure is not drawn to scale.

Enzymatic modifications

- **Phosphorylation/dephosphorylation**
 - On serine, threonine, tyrosine
- **Glycosylation**
 - N-glycosylation (asparagine-linked)
 - O-glycosylation (serine-, threonine-linked)
- **N-Acetylation/deacetylation**
 - On lysines
- **N-Methylation/demethylation**
 - On lysines - mono-, di- and trimethylation


Steve Barnes BMG 744 2-8-08

Phosphorylation of proteins

- In some cases, proteins are normally found in a stable, hyperphosphorylated state, e.g., casein
- In many cases, it is a transient event that causes 10-100 fold increase in enzyme activity. This is the way signals are propagated through a signal transduction pathway. However, the molar abundance of phosphorylation at an individual site may only be 1-2%.

Steve Barnes BMG 744 2-8-08

Chemistry of phosphorylation



Steve Barnes BMG 744 2-8-08

Mass spectrometry of phosphorylated proteins

- Adds H₃PO₄ (+98)
- Eliminates water (-18)
- Net change +80
- if the phosphate ion is released (i.e., in negative ion spectra), it is seen as *m/z* 79 and/or *m/z* 63
- Phosphate can be confused with sulfate and bromide (sulfate, 79.9568 Da; phosphate, 79.9663 Da; and ⁸¹Br, 79.9083 Da)

Steve Barnes BMG 744 2-8-08

Limitations of mass spectrometry

- Although it can deliver sensitivity in the low fmol range (similar to immunological methods), because it is a universal detection method, finding the needle in the “haystack” of all the other peptides is a challenge
- Recovering the phosphopeptides from the matrix of the sample is more important than the mass spectrometry measurement

Steve Barnes BMG 744 2-8-08

Finding a phosphate group

Several methods are in current use for detection of phosphopeptides

- use of parent ion or neutral loss scanning
- phosphatase sensitivity
- affinity methods for enrichment of phosphopeptides
 - anti-phospho-Ser/Thr/Tyr antibodies
 - metal ion affinity
 - chemical reaction/biotin affinity



Steve Barnes BMG 744 2-8-08

Parent ion scanning to detect phosphopeptides

- The procedure depends on the detection of the m/z 79 ion fragment (PO_3^-) during collision-induced dissociation in a triple quadrupole instrument operating in the negative ion mode
- *Parent ion scanning* is a reversal of the more familiar daughter ion MS-MS where the parent ion is selected (in Q1) and a mass spectrum of the daughter ion fragments is obtained by scanning in Q3
- In *parent ion scanning*, the daughter ion fragment (in this case m/z 79) is held constant in Q3 and a mass spectrum of parent ions that give rise to the daughter ion obtained by scanning in Q1.
- Having identified the phosphopeptides, the sample can be re-analyzed to obtain daughter ion MS-MS spectra on selected ions in the positive ion mode



Steve Barnes BMG 744 2-8-08

Parent ion scan to detect phosphopeptide




Steve Barnes BMG 744 2-8-08 [Annan Anal Chem 73:393](#)

Unknown casein phosphopeptide



Steve Barnes BMG 744 2-8-08 [Annan Anal Chem 73:393](#)

Deducing phosphopeptide sequence



Steve Barnes BMG 744 2-8-08 [Annan Anal Chem 73:393](#)

Recovery and enhancement of phosphopeptides

The biggest problem in the detection of phosphopeptides is how to convert the initial sample matrix into a form suitable for mass spectrometry analysis.

- how to handle minute samples with minimal losses
- how to recover and detect all the phosphopeptides
- how to recover and detect the non-phosphorylated proteins to determine the extent of phosphorylation at individual sites

Steve Barnes BMG 744 2-8-08


Antibodies and phosphopeptides

- In this approach, both the phosphorylated and non-phosphorylated forms of a protein may be recovered from the sample matrix
- This can most easily be achieved by immunoprecipitation of the protein with an antibody that recognizes epitope(s) that is(are) in common with both forms


NOTE *that as with all immunoprecipitation methods, the best results will be achieved if the antibody is coupled to agarose beads. This allows selective immuno-absorption of the antigen, washing free of contaminating proteins, AND following elution, minimization of the amount of antibody that is in the eluate and therefore would be analyzed by mass spectrometry*

Steve Barnes BMG 744 2-8-08


Variability of anti-phosphoserine and anti-phosphothreonine antibodies



Steve Barnes BMG 744 2-8-08 [Gronborg et al., 2002](#)



Phosphorylation in HeLa cells stimulated by calyculin



Western blot analysis with anti-phosphoserine and anti-phosphotyrosine antibodies

Steve Barnes BMG 744 2-8-08 [Gronborg et al., 2002](#)

Detection of phosphopeptides based on their sensitivity to phosphatase

- An alternative source of potentially phosphorylated proteins are individual spots on 2D-IEF/SDS gels. The protein preparation so isolated is either hydrolyzed by trypsin in solution (or in the gel piece) or using solid-phase trypsin
- One portion of the resulting tryptic peptides (in 50% acetonitrile:water) is analyzed by MALDI-TOF-MS. A second portion is diluted into 50 mM NH_4HCO_3 buffer and reacted with 0.5 U calf intestinal alkaline phosphatase at 37°C for 30 min. Sample is dried with a SpeedVac, redissolved in 50% acetonitrile:water, and re-analyzed by MALDI-TOF-MS


Steve Barnes BMG 744 2-8-08

Identifying phosphopeptides using alkaline phosphatase

- Phosphopeptides shift down by m/z 80 (or units of 80 in the case of multiply phosphorylated peptides)
- The peaks identified as phosphopeptides can then be analyzed in a nanoelectrospray experiment where collision-induced dissociation is used to determine the identity of the peptide and the phosphorylation site in the sequence

Steve Barnes BMG 744 2-8-08

Detecting a phosphopeptide with alkaline phosphatase



Steve Barnes BMG 744 2-8-08


Zhang et al., 2000
MS in Biology & Medicine

Selective enhancement of phosphopeptides in tryptic digests

- Immobilized metal affinity chromatography (IMAC). Similar to Ni-affinity resins used in the purification of 6xHis-tagged proteins. The affinity phase can be charged with different metal ions (as their chlorides)
- Fe(III) and Ga(III), and to a lesser extent Zr(IV), were the most effective for the recovery of two synthetic phosphopeptides
- A tryptic digest containing both phosphorylated and non-phosphorylated peptides is passed over the IMAC column at acid pH (pH 2.5-3)
- The column is washed with 0.1 M acetic acid to remove unbound peptides
- Elute with sodium phosphate (have to desalt) or with NH_4OH
- Esterification may prevent Asp- or Glu-containing peptides from binding

Steve Barnes BMG 744 2-8-08

Selective biotinylation of phospho-groups



Goshe et al., Anal Chem 73: 2578 (2001)
Adamcyk et al., Rapid Commun Mass Spec 15; 1481 (2001)

Steve Barnes BMG 744 2-8-08

How to identify phosphorylated peaks by searching databases

- The databases you have used earlier in this class have some ability to predict the expected masses for a limited number of posttranslational modifications
 - MASCOT (<http://www.matrixscience.com>)
 - PROTEIN PROSPECTOR (<http://prospector.ucsf.edu/>)

Steve Barnes BMG 744 2-8-08

How to identify posttranslational modifications

FindMod at

<http://www.expasy.org/tools/findmod/>

It examines mass fingerprinting data for mass differences between empirical and theoretical peptides. If the mass difference corresponds to a known modification, it also makes intelligent guesses as to the site of modification.

Steve Barnes BMG 744 2-8-08

FindMod

Some of the modifications considered are:

acetylation	amidation	biotinylation
C-mannosylation	deamidation	flavinilylation
farnesylation	formylation	geranyl-geranylation
γ -carboxyglutamic acid	hydroxylation	lipoylation
methylation	myristoylation	N-acyl diglyceride
O-GlcNac	palmitoylation	phosphorylation
pyridoxal phosphate	phospho-pantetheine	pyrrolidone-carboxylic acid
sulfation		

NOTE that none of the common chemical modifications (alkylation of sulfhydryl groups with iodoacetic acid, iodoacetamide, 4-vinylpyridine, and acrylamide) were included.

Wilkins M.R., Gasteiger E., Gooley A., Herbert B., Molloy M.P., Binz P.A., Ou K., Sanchez J.-C., Bairoch A., Williams K.L., Hochstrasser D.F. *High-throughput Mass Spectrometric Discovery of Protein Post-translational Modifications*. [Journal of Molecular Biology, 289, p. 645-657 \(1999\)](#)
Gasteiger E., Hoogland C., Gattiker A., Duvaud S., Wilkins M.R., Appel R.D., Bairoch A.; *Protein Identification and Analysis Tools on the ExPASy Server*; (In) John M. Walker (ed): [The Proteomics Protocols Handbook. Humana Press \(2005\)](#).

Steve Barnes BMG 744 2-8-08

Site for compilation of PTMs

<https://www.abrf.org/index.cfm/dm.home>

This site was put together by Ken Mitchelhill, Len Packman and friends

Currently ranges from dephospho (-79) to (Hex)3-HexNAc-(dHex)HexNAc (+1,039)

Steve Barnes BMG 744 2-8-08

QuickTime™ and a decompressor are needed to see this picture.

From Nature Methods 4:787 (2007)

Steve Barnes BMG 744 2-8-08

QuickTime™ and a
decompressor
are needed to see this picture.

From Nature Methods 4:787 (2007)

Steve Barnes BMG 744 2-8-08

QuickTime™ and a
decompressor
are needed to see this picture.

From Nature Methods 4:787 (2007)

Steve Barnes BMG 744 2-8-08

Detecting PTMs

- If we don't have any idea about what the modification(s) is(are), how do we proceed?
- We won't have the modification on MASCOT or Protein Prospector
- No antibodies or convenient affinity phases


Steve Barnes BMG 744 2-8-08

Can we approach this globally?

- It's asking a lot
 - Too many degrees of freedom
 - Endless modifications
- Should we try?
 - John Yates' group has
 - [MacCoss et al. PNAS 99:7900 \(2002\)](#)

Steve Barnes BMG 744 2-8-08

MudPIT - Multi-dimensional Protein Identification Technology



Steve Barnes BMG 744 2-8-08



John Yates

The Yates approach - selection of “proteomes”

- BioRad mixed molecular weight standards (10 pmol each) plus 1 pmol of a phosphorylated glycogen phosphorylase
- Cdc2 protein complex isolated with TAP (15 μ g)
- Lens proteins from 4-yr old

Steve Barnes BMG 744 2-8-08

MacCoss et al, 2002



Hydrolysis procedures

Reduced, carboxymethylated in 8 M urea

Diluted x 3 with
100 mM Tris-HCl,
pH 8.5

100 mM CaCl₂
added to 1 mM

Roche **trypsin** 1:50,
incubated 12-24 h
at 37°C

quenched with
90% formic acid
to 4%

Diluted x 3 with 4.8
M Urea-100 mM Tris-
HCl, pH 8.5

subtilisin added
1:50, incubated 2-3
hr at 37°C

quenched with
90% formic acid
to 4%

Steve Barnes BMG 744 2-8-08


Diluted x 3 with
100 mM Tris-HCl,
pH 8.5

incubated with
elastase 1:50 12 h
at 37°C

quenched with
90% formic acid
to 4%

MacCoss et al, 2002

Elution from a triphasic column




Steve Barnes BMG 744 2-8-08

MacCoss et al, 2002

Automated MS-MS analysis

- Limit analysis to 2⁺ or 3⁺-charged peptides
- Delete poor quality spectra
- Identify peptides with 98-Da neutral loss
- Analyze the remaining spectra with SEQUEST (operating on a 31-node beowulf computer cluster) to identify proteins
- Search sequences of identified proteins for PTMs by 80 (STY phosphorylation), 42 (K acetylation), 16 (MWY oxidation) and 14 (K methylation)

Steve Barnes BMG 744 2-8-08 [MacCoss et al, 2002](#)



Steve Barnes BMG 744 2-8-08

Results

Protein standard mixture:

- The three digests were combined - 83.7% and 95.4% coverage for glycogen phosphorylase and BSA
- Identified the glycogen phosphorylase phosphorylation site as well as the two known sites (S69/S345) in ovalbumin
- N-acetylation found plus many sites of methionine oxidation (? due to work up or real)
- New sites found - phosphorylation at S237/S241 in ovalbumin and methylation at R652

Steve Barnes BMG 744 2-8-08 [MacCoss et al, 2002](#)

Results

Cdc2p complex:

- 200 proteins - 20 showed >40% sequence coverage
- Expected Y15 and T167 phosphorylation sites on Cdc2p found
- New phosphorylation sites found on cyclin partners Cdc13 and Cig1p
- Multiple methylation sites on Cdc2p

Steve Barnes BMG 744 2-8-08 [MacCoss et al, 2002](#)

Results

Lens proteins:

- Found 272 proteins - 52 had >40% sequence coverage
- 90% are crystallins
- PTMs accumulate over your lifetime
- Used 18-step MudPIT because of complexity
- 73 different PTMs found on the 11 crystallins
- Found the 13 of 18 PTMs previously described in all species
- Found 60 other new PTMs in phosphorylation, oxidation, acetylation and methylation

Steve Barnes BMG 744 2-8-08 [MacCoss et al, 2002](#)

Quantitative proteomics

Use of isotopes

- ICAT (d_0/d_8) and ICAT $^{13}C_0/^{13}C_8$
- d_0/d_{10} propionic anhydride (N-terminal labeling)
- $^{15}N/^{14}N$ (whole cell labeling)
- $^{18}O/^{16}O$ (trypsin)
- iTRAQ labeling
- Non-isotope methods
 - Peptide coverage
 - Classical triple quadrupole methods

Steve Barnes BMG 744 2-8-08

Quantification methods
in mass spectrometry

From review in Nature
Methods 4-79 (2007)

QuickTime™ and a
decompressor
are needed to see this picture.


Steve Barnes BMG 744 2-8-08

Isotope-coded affinity technology

Isotope-Coded Affinity Tags

heavy reagent: D8-ICAT Reagent (X=deuterium)



light reagent: D0-ICAT Reagent (X=hydrogen)



This reagent reacts with cysteine-containing proteins (80-85% of proteome)


Labeling can be replacement of hydrogens (X) with deuterium, or better to
exchange ^{12}C with ^{13}C in the linker region (this avoids chromatography issues)

Steve Barnes BMG 744 2-8-08




SILAC

- **SILAC, stable isotope labeling by amino acids in cell culture, is being used to quantify proteins.**



Steve Barnes BMG 744 2-8-08 [Ong et al., MCP 1:367, 2002](#)

Time-dependent leucine incorporation with SILAC




The cells are pre-labeled with leucine-d₀. Leucine-d₃ is added to the medium and cells sampled at various times later. The peaks annotated with d0 and d3 are the triply charged peaks of the peptide VAPEEHPVLLTEAPLNPK, which contains three leucines.

Steve Barnes BMG 744 2-8-08 [Ong et al., MCP 1:367, 2002](#)

¹⁸O-labeling

- Trypsin catalyzes the transfer of ¹⁸O in ¹⁸O-enriched water to both the carboxylate oxygens of the C-terminus of tryptic peptides




- The peptides have an increase in mass of 4 Da
- Generally not considered a large enough mass difference

Steve Barnes BMG 744 2-8-08



iTRAQ quantification

- The iTRAQ™ reagents
 - React with Lys amino groups and each one adds 145 Da to the molecular weight of the peptide
 - Fragmentation produces reporter ions from *m/z* 114, 115, 116 and 117
 - New iTRAQ kit contains 8 forms with reporter fragment ions of *m/z* 114, 115, 116, 117, 118, 119 and 121



Steve Barnes BMG 744 2-8-08


iTRAQ™ Reagent Design



Other non-isotopic quantitative methods in proteomics

The coverage (the number of peptides observed for a protein) is sensitive to the amount of the protein

- This can be used to calculate whether a treatment affects the abundance of a protein where fold-change > 2
- Applies to LC-MS (MUDPIT methods)




Steve Barnes BMG 744 2-8-08

Triple quad MRM analysis

Peptides of interest can be analyzed like small molecules

- Choose the parent molecular ion, collide with argon gas and select a unique fragment



- **Multiple reaction ion scanning**


First filter the [M-H]⁻ molecular ion of the analyte (Q1)

Fragment the molecular ion with N₂ gas (Q2)

Select a specific (and unique) fragment ion (Q3)



Steve Barnes BMG 744 2-8-08



Quantitation experiment for biotinylated cytochrome c MRM analysis monitored in 50 channels



Each colored peak represents a different biotinylated peptide

Steve Barnes BMG 744 2-8-08





Application of LC-MRM-MS to the plasma proteome

Identify the proteins of interest


In silico, generate the tryptic peptides from each protein

Select the best combination of parent peptide and fragment y ion

Determine the expected y ions for each peptide and compare to y ions of all other tryptic peptides of known human proteins that have masses within $\pm 1\text{Da}$

Steve Barnes BMG 744 2-8-08

Slick way of making ^{13}C -labeled peptide internal standards



Quantitative peptide MRM-MS

- The albumin-depleted plasma proteome is mixed with the composite ^{13}C , ^{15}N -labeled protein internal standard and then treated with trypsin
- The molecular ions (doubly charged) and the specific y ions for each peptide and its labeled form are entered into the MRM script one channel at a time
- A single run may consist of 30 peptides in 60 channels
- Sensitivity is compromised by “sharing out” measurement time, but can be compensated for by carrying out nanoLC

Steve Barnes BMG 744 2-8-08

Advantage of a C-terminal labeled lysine

186 301 448 505 642 755 886 987 1115 b ions
A D E F G H I M T K
1133 1062 948 833 686 629 492 379 248 147 y ions

With the labeled lysine at the C-terminus, only the b_{10} ion contains the isotope atoms

186 301 448 505 642 755 886 987 1123 b ions
A D E F G H I M T K*
1141 1070 956 841 694 637 500 387 256 155 y ions

Steve Barnes BMG 744 2-8-08

References for this talk (1)

- Annan RS, Hudleston MJ, Verma R, Deshaies RJ, Carr SA. A multidimensional electrospray MS-based approach to phosphopeptide mapping. *Anal. Chem.* 73:393, 2001.
- Flory MR, Griffin TJ, Martin D, Aebersold R. Advances in quantitative proteomics using stable isotope tags. *Trends in Biotechnology* 20: S23, 2002.
- Taupenot L, Harper KL, O'Connor DT. The chromogranin-secretogranin family. *New Engl. J. Med.* 348: 1134, 2003.
- Lam YW, Mobley JA, Evans JE, Carmody JF, Ho S-M. Mass profiling-directed isolation and identification of a stage-specific serologic protein biomarker of advanced prostate cancer. *Proteomics* 5: 2927, 2005.

Steve Barnes BMG 744 2-8-08

Bibliography (2)

- Ong SE, Mann M. Mass spectrometry-based proteomics turns quantitative. *Nature Chemical Biology.* 1:252-262, 2005.
- Gruhler A, Schulze WX, Matthiesen R, Mann M, Jensen ON. Stable isotope labeling of *Arabidopsis thaliana* cells and quantitative proteomics by mass spectrometry. *Molecular & Cellular Proteomics.* 4:1697-1709, 2005.
- Anderson L, Hunter CL. Quantitative Mass Spectrometric Multiple Reaction Monitoring Assays for Major Plasma Proteins. *Molecular & Cellular Proteomics* 5:573-588, 2006.
- Yao X, Freas A, Ramirez J, Demirev PA, Fenselau C. Proteolytic ¹⁸O labeling for comparative proteomics: model studies with two serotypes of adenovirus. *Analytical Chemistry* 73, 2836-42, 2001.
- Wang G, Wu WW, Zeng W, Chou C-L, and Shen R-F. Label-Free Protein Quantification Using LC-Coupled Ion Trap or FT Mass Spectrometry: Reproducibility, Linearity, and Application with Complex Proteomes. *Journal of Proteome Research* 5: 1214-1223, 2006.

Steve Barnes BMG 744 2-8-08