

Ion fragmentation of small molecules in mass spectrometry

Jeevan Prasain

jprasain@uab.edu

6-2612

Feb 3, 2012


Small molecules are important!!

- 89% of all known drugs and 50% of all drugs are derived from pre-existing metabolites.
- Small molecules are cofactors and signalling molecules to 1000's of proteins.
- 100,000 (lipidome)




Feb 3, 2012

Triglycerides




Genistein
(a plant secondary metabolite)



Taxol


Nomenclature: the main names and acronyms used in mass spectrometry

- **Molecular ion:** Ion formed by addition or the removal of one or several electrons to or from the sample molecules-**Electron Impact (EI-MS)**. $M + e^- \rightarrow M^{+\bullet} + 2e^-$
- **Adduct Ion:** Ion formed through interaction of two species and containing all the atoms of one of them plus one or several atoms of them (e.g. alkali, ammonium).



Feb 3, 2012


Adduct formation in +/-ve ion modes



Feb 3, 2012

Nielsen et al., J Nat Prod. 2011

Molecules with inherent positive charge- molecular weight and m/z are same



Feb 3, 2012

Contd..

- **Pseudomolecular ion:** Ion originating from the analyte molecule by abstraction of a proton $[M-H]^-$ or addition of proton $[M+H]^+$
- **Tandem mass spectrometry (Cooks, 1976): MS/MS (McLafferty, 1978),** tandem in space or time
- **Precursor ion/parent ion:** Ions undergoing fragmentation.
- **Product ion/daughter ion:** Ions resulting from parent/precursor ions.
- **Neutral loss:** Fragments lost as neutral molecules
- **In positive ionization mode,** a trace of formic acid is often added to aid protonation of the sample molecules; in **negative ionization mode** a trace of ammonia solution or a volatile amine is added to aid deprotonation of the sample molecules. Proteins and peptides are usually analysed under positive ionization conditions and polyphenols and acids under negative ionization conditions. In all cases, the m/z scale must be calibrated.


Feb 3, 2012

Isotopic distribution and MS

- $1\text{H} = 99.9\%$, $2\text{H} = 0.02\%$
- $12\text{C} = 98.9\%$, $13\text{C} = 1.1\%$
- $35\text{Cl} = 68.1\%$, $37\text{Cl} = 31.9\%$
- **Monoisotopic mass** - the mass of the most abundant isotope
- **Average mass**- the abundance weighted mass of all isotopic components.

Feb 3, 2012

What is Collision Induced Dissociation (CID) or Collisionally Activated Dissociation (CAD) ?



Schematic of CID fragmentation

Other activation processes:


PSD (post source-decay)

ECD (electron capture dissociation)

SID (surface-induced dissociation)

Feb 3, 2012

Various types of MS/MS experiments



Mode of operation	Q1	q2	Q3
Q1 Scan	Resolving (Scan)	RF-only	RF-only
Q3 Scan	RF-only	RF-only	Resolving (Scan)
Product Ion Scan (PI)	Resolving (Fixed)	Fragment	Resolving (Scan)
Precursor Ion Scan (PC)	Resolving (Scan)	Fragment	Resolving (Fixed)
Neutral Loss Scan (NL)	Resolving (Scan)	Fragment	Resolving (Scan Offset)
Selected Reaction Monitoring mode (SRM)	Resolving (Fixed)	Fragment	Resolving (Fixed)

Enhanced Q3 Single MS (EMS)	RF-only	No frag	Trap/scan
Enhanced Product Ion (EPI)	Resolving (Fixed)	Fragment	Trap/scan
MS ²	Resolving (Fixed)	Fragment	Isolation/frag trap/scan
Time delayed fragmentation (TDF)	Resolving (Fixed)	Trap/No frag	Frag/trap/scan
Enhanced Resolution Q3 Single MS (ER)	RF-only	No frag	Trap/scan
Enhanced Multiply Charged (EMC)	RF-only	No frag	Trap/scan

Figure 1. Schematic of QqLIT (Q TRAP, AB/MDS, Sciex) and description of the various triple quadrupole and trap operation modes.

Hopfgartner et al. J. Mass Spectrom, 2004

Feb 3, 2012

Applications of MS/MS


- **Pharmaceuticals**- Identification and quantification of drug metabolites, PK/PD
- **Academic/biotechnology**- analysis of protein/peptides, authentication and profiling of chemical components in a crude mixture, substructure analysis of unknown components
- **Clinical**- eg. neonatal screening, steroids in athletes etc.
- **Environment**- eg. dioxins in fish..
- **Geological**- eg. oil compositions...

Feb 3, 2012

Interpreting MS/MS spectra

- Likely sites of protonation or deprotonation.
- Likely leaving group.
- Literature study


Where are the sites of deprotonation/protonation?
What is the most likely leaving group in this molecule?



Fragmentation always follows the basic rules of chemistry

Feb 3, 2012

O- and C-glucosides fragment differently in ESI-MS/MS




Puerarin
 m/z 415

Feb 3, 2012


Prasain et al. J. Agric. Food Chem. 2003

Possible product ions of puerarin m/z 415 in MS/MS



Feb 3, 2012

Intensity of product ions indicates their stability - ions bearing aromatic ring are more intense




Ion fragmentation for identification of phase II drug metabolites (glucuronide/sulfate conjugates)

Feb 3, 2012

What fragment ions are characteristic for glucuronide conjugates?


Product ion spectrum of genistein glucuronide in ESI-MS/MS




Glucosides/glucuronides conjugates are easily cleaved off by higher potential at orifice

Feb 3, 2012


MSMS of m/z 429 indicate that it may be daidzein glucuronide



The loss of 80 Da from the parent ion and the presence of m/z 80 in the product ion spectra are the indicative of sulfate conjugates of like daidzein [A] and equol [B]




What happens with aliphatic sulfates in MS/MS?



Aliphatic and aromatic sulfate conjugates behave differently in MS/MS, aliphatic typically show m/z 97 (HSO_4^-) and m/z 80 (SO_3^-)

Feb 3, 2012

The absence of the m/z 97 fragment with the base peak m/z 80 makes the distinction between aromatic and aliphatic sulfates



Feb 3, 2012 Source: Weidolf et al. Biomed. and Environ. Mass Spec. 1988

Change in mass is associated with possible metabolic reaction

Metabolic rxn	Change in mass
Methylation	14
Demethylation	-14
Hydroxylation	16
Acetylation	42
Epoxidation	16
Desulfuration	-32
Decarboxylation	-44
Hydration	18
Dehydration	-18

Feb 3, 2012

Characteristic fragmentation of drug conjugates by MS/MS


Conjugate	Ionization mode	Scan
Glucuronides	pos/neg	NL 176 amu
Hexose sugar	pos/neg	NL 162 amu
Pentose sugar	pos/neg	NL 132 amu
Phenolic sulphate	pos	NL 80 amu
Phosphate	neg	Precursor of m/z 79
Aryl-GSH	pos	NL 275 amu
Aliphatic-GSH	pos	NL 129
taurines	Pos	Precursor of m/z 126
N-acetylcysteins	neg	NL 129 amu

Feb 3, 2012

NL = neutral loss.


Kostiainen et al., 2003

Analysis of steroids by MS/MS




Feb 3, 2012

Estradiol Standard Curve 0.05 – 25 μM $r = 0.9959$



Feb 3, 2012


Derivatization of estradiol with dansyl chloride leads to the formation of E₂-dansyl (m/z 506)



Source: Nelson et al. Clinical Chemistry, 2004


Feb 3, 2012

Derivatization tremendously helps increase sensitivity of E₂



Feb 3, 2012

**Calibration curve for dansylated E2 showing
linearity from 0.005-100 nM concentration range
($r = 0.999$)**




Feb 3, 2012

**Substructure analysis in ESI-MS/MS
(dereplication and partial identification
of natural products)**


Feb 3, 2012

Fragmentation of basic taxoids from *T. Wallichiana* extract


Prasain et al. Anal Chem, 2001



[1] MW=861




[2] MW=769



[3] MW=683


Feb 3, 2012

ESI-MS/MS spectra of taxoids (1-3). Peaks m/z 194 and 210 represent the intact alkaloid side chain.




Feb 3, 2012

MS/MS precursor-scan spectra of typical alkaloid side chains to identify the basic taxoids compounds in an ethyl acetate extract of *T. wallichiana*.



Feb 3, 2012

Comparison of precursor scan spectra obtained from the scaffold m/z 309 and side chain m/z 194, 210 and 252



Taxoids with scaffold m/z 309 and alkaloid side chains are shown by dashed lines

Feb 3, 2012

References

1. **Electrospray Ionization Mass Spectrometry** by Richard B. Cole.
2. **Stefanowicz P, Prasain JK, Yeboah KF, Konishi Y. Detection and partial structure elucidation of basic taxoids from *Taxus wallichiana* by electrospray ionization tandem mass spectrometry. Anal Chem. 2001;73:3583-9.**
3. [Prasain J.K., Wang C.-C., Barnes S. Mass spectrometric analysis of flavonoids in biological samples. *Free Radical Biology & Medicine*, 37: 1324-1350, 2004.](#)
4. **William Griffiths. Tandem mass spectrometry in the study of fatty acids, bile acids and steroids. Mass Spectrometry Reviews, 2003;22:81-152.**
5. **Yi et al., Anal Bioanal Chem. 2006.**

Feb 3, 2012