

#### **Dissociative research**

- Samples are collected and stored for analysis at a "later" time
- "Later" can be months or years after sample collection
  - Of little direct benefit to the patient
  - Although may influence the community of patients
  - True of many analyses

#### **Real time analysis**


- Existing, familiar applications
- Gases!
- The iknife
  - GI surgery
  - Cancer margins
  - Pathology
- DESI
- CARS

#### **Real-time analysis**

- We see the real-time use of MS when we go through security checks at the airport
  - Checks for ion signatures of explosives
- Other devices are used to check for specific volatiles in the breath







#### **Gases produced in the GI tract**

- H<sub>2</sub>, CO<sub>2</sub> and CH<sub>4</sub> from carbohydrates
  - Firmicutes
  - From pyruvate and NAD(P)H/FADH<sub>2</sub>
  - H<sub>2</sub> used by sulfate-reducing bacteria (SRBs), methanogenic Archaea, and acetogens
- SRBs produce H<sub>2</sub>S
- NO from nitrates

#### Methods for measuring gases


Technology	Operation mode	Target intestinal gas	Detection limit	Cross-sensitivity	Response time	Life time	Estimated cost
Spectrometry bas	ed <sup>a</sup>				4.000		
GC-MS	Off line	All gases	ppt to ppb	Low	~Several minutes	Long	>US\$300k
IMS	Real time	All gases	ppb	Low	<1 min	Long	>US\$100k
PTR-MS	Real time	All gases	ppt	Low	<1 min	Long	>US\$400k
SIFT-MS	Real time	All gases	ppb	Low	<1 min	Long	>US\$400k
LS	Real time	Most gases except H <sub>2</sub>	ppt to ppb	Low	<1 min	Long	<us\$50k< td=""></us\$50k<>
Sensor based							
Electrochemical	Real time	H <sub>2</sub> , H <sub>2</sub> S, NO, and CO <sub>2</sub>	ppm	Medium	<30 s	Short	<us\$100< td=""></us\$100<>
Calorimetric	Real time	H <sub>2</sub> , CH <sub>4</sub> , and CO <sub>2</sub>	ppt	High	<10 s	Medium	<us\$100< td=""></us\$100<>
NDIR	Real time	CO2, CH4, and VOCs	ppm to ppt	Low	<20 s	Long	<us\$300< td=""></us\$300<>


GC-MS gas chromatography-mass spectrometry
IMS ion mobility mass spectrometry

PTR-MS proton transfer reaction mass spectrometry SIFT-MS selection ion flow tube-mass spectrometry


LS laser spectrometry

Jian Zhen Ou et al., Trends Biotech, 2015






#### Real-time in situ monitoring gas production




- The device is swallowed
- Completes full mouth-toanus transit, reporting data as it goes
- Also provides positional information
- Operates at 405, 433, and 915 MHz
- Uses Lithium batteries!!

Jian Zhen Ou et al., Trends Biotech, 2015



### **The Challenge for Mass Spec**



How to get the mammoth into the gas phase for analysis?

### **Droplet principle of electrospray**




#### **Droplet spray**

- Sneeze
- Lung motion
- Surgical knife
- Other vapors










### **Link to videos by James Kinross**

Colorectal surgeon from Imperial College, London Plenary Speaker at the UAB 2016 Metabolomics Workshop


http://www.uab.edu/proteomics/metabolomics/workshop/2016/videos/kinross\_day 2.html


 $\underline{\text{http://www.uab.edu/proteomics/metabolomics/workshop/2016/videos/kinross2}} \ \ \underline{\text{day}} \\ \underline{\text{2.html}}$ 




## Phospholipid patterns are characteristic of cells and tissues

- · Single items are not sufficient as biomarkers
- The classes of phospholipids and their fatty acid composition contain pattern discriminators
- In the absence of known classifiers, principal components analysis looks for groups of components that have the larger sources of variation
  - An individual sample's contributions to these groups are plotted in a 2D or 3D manner








Computer-driven, Rapid Evaporative Imaging MS (REIMS) for tissue sections






Golf et al., Anal Chem 2015

#### **Modes of data acquisition for REIMS**


Line Scans: Cutting Mode




Individual Pixels: Pointing Mode




Golf et al., Anal Chem 2015






#### **Desorption electrospray ionization (DESI)**


 Works by directing an electrical fine spray at a tissue target – does not require deposition of a matrix





## The IDH story of brain and other tumors


- IDH1 (isocitrate dehydrogenase) is mutated in position
   132 in a GWAS study of patients with glioblastomas
- IDH1 catalyzes the conversion of isocitrate to alphaketoglutarate ( $\alpha$ KG) which is a two-step reaction
- Mutant IDH1 catalyzes the first step to 2-hydroxyglutarate (2HG), but not the second one to  $\alpha$ KG
- 2HG is considered to be an onco-metabolite
- What follows is a study from a group at Harvard –
  performed in the Advanced Multimodality Image Guided
  Operating Suite at Brigham and Women's Hospital
































Distribution of glucose/citrate ratio of some representative prostate tissue specimens showing significant elevation of the glucose/citrate ratio in cancer. The Top of each panel (A–O) shows the histopathological evaluation (H&E) of the corresponding tissue, where cancer areas have been demarcated by red, benign areas by black, stroma areas by green, and inflammation areas by blue.

## **Use of Raman spectroscopy**Real-time imaging of metabolites in skin

• <a href="http://bernstein.harvard.edu/research/cars-why.htm">http://bernstein.harvard.edu/research/cars-why.htm</a>



Sunny Xie, PhD - Harvard

# The future of medicine and surgery

http://www1.imperial.ac.uk/phenomecentre/

#### **Publications**

- Santagata S, Eberlin LS, Norton I, Calligaris D, Feldman DR, Ide JL, Liu X, Wiley JS, Vestal ML, Ramkissoon SH, Orringer DA, Gill KK, Dunn IF, Dias-Santagata D, Ligon KL, Jolesz FA, Golby AJ, Cooks RG, Agar NY. <u>Intraoperative mass spectrometry mapping of an oncometabolite to guide brain tumor surgery.</u> PNAS 2014;111(30):11121-6.
- Banerjee S, Zarea RN, Tibshirani RJ, Kunder CA, Nolley R, Fan R, Brooks JD, Sonn GA.
   Diagnosis of prostate cancer by desorption electrospray ionization mass spectrometric imaging of small metabolites and lipids. PNAS early edition, March 2017
- Golf O, Strittmatter N, Karancsi T, Pringle SD, Speller AV, Mroz A, Kinross JM, Abbassi-Ghadi N, Jones EA, Takats Z. Rapid evaporative ionization mass spectrometry imaging platform for direct mapping from bulk tissue and bacterial growth media. <a href="https://doi.org/10.1007/j.chm">Anal Chem. 2015 Mar 3;87(5):2527-34</a>.
- In vivo endoscopic tissue identification by rapid evaporative ionization mass spectrometry (REIMS).Balog J, Kumar S, Alexander J, Golf O, Huang J, Wiggins T, Abbassi-Ghadi N, Enyedi A, Kacska S, Kinross J, Hanna GB, Nicholson JK, Takats Z. <u>Angew Chem Int Ed Engl. 2015 Sep 14;54(38):11059-62</u>.