



GBSC 724 March 20, 2023

Real-time connection of Metabolomics with Medicine and Surgery and the rest of life

Stephen Barnes, PhD, FASN
Professor of Pharmacology & Toxicology
Director, Targeted Metabolomics and
Proteomics Laboratory





Dissociative research

- Samples are collected and stored for analysis at a "later" time
- "Later" can be months or years after sample collection
 - Of little direct benefit to the patient
 - Although may influence the community of patients
 - True of many analyses

3

Link to videos by James Kinross

Colorectal surgeon from Imperial College, London Plenary Speaker at the UAB 2016 Metabolomics Workshop

http://www.uab.edu/proteomics/metabolomics/workshop/2016/videos/kinross day2.html

http://www.uab.edu/proteomics/metabolomics/workshop/2016/videos/kinross2 day2.html

Real time analysis

- Existing, familiar applications
- Gases!
- The iknife
 - GI surgery
 - Cancer margins
 - Pathology
- DESI
- CARS
- Raman

5

Real-time analysis

- We see the real-time use of MS when we go through security checks at the airport
 - Checks for ion signatures of explosives
- Other devices are used to check for specific volatiles in the breath





Noses and smell – real time analysis



The superior volatile metabolite detector

7

Gases produced in the GI tract

- H₂, CO₂ and CH₄ from carbohydrates
 - Firmicutes
 - From pyruvate and NAD(P)H/FADH₂
 - H₂ used by sulfate-reducing bacteria (SRBs), methanogenic
 Archaea, and acetogens
- SRBs produce H₂S
- NO from nitrates

Methods for measuring gases

Technology	Operation mode	Target intestinal gas	Detection limit	Cross-sensitivity	Response time	Life time	Estimated cost
Spectrometry bas	The state of the s				time		COST
GC-MS	Off line	All gases	ppt to ppb	Low	~Several minutes	Long	>US\$300k
IMS	Real time	All gases	ppb	Low	<1 min	Long	>US\$100k
PTR-MS	Real time	All gases	ppt	Low	<1 min	Long	>US\$400k
SIFT-MS	Real time	All gases	ppb	Low	<1 min	Long	>US\$400k
LS	Real time	Most gases except H ₂	ppt to ppb	Low	<1 min	Long	<us\$50k< td=""></us\$50k<>
Sensor based ^b							
Electrochemical	Real time	H ₂ , H ₂ S, NO, and CO ₂	ppm	Medium	<30 s	Short	<us\$100< td=""></us\$100<>
Calorimetric	Real time	H ₂ , CH ₄ , and CO ₂	ppt	High	<10 s	Medium	<us\$100< td=""></us\$100<>
NDIR	Real time	CO ₂ , CH ₄ , and VOCs	ppm to ppt	Low	<20 s	Long	<us\$300< td=""></us\$300<>

GC-MS gas chromatography-mass spectrometry

IMS ion mobility mass spectrometry


PTR-MS proton transfer reaction mass spectrometry SIFT-MS selection ion flow tube-mass spectrometry

LS laser spectrometry


Jian Zhen Ou et al., Trends Biotech, 2015


9


Device for measuring fecal gas production




Jian Zhen Ou et al., Trends Biotech, 2015









Droplet principle of electrospray





Droplet spray

- Sneeze
- Lung motion
- Surgical knife
- Other vapors












Τ/


Phospholipid patterns are characteristic of cells and tissues

- Single items are not sufficient as biomarkers
- The classes of phospholipids and their fatty acid composition contain pattern discriminators
- In the absence of known classifiers, principal components analysis looks for groups of components that have the larger sources of variation
 - An individual sample's contributions to these groups are plotted in a 2D or 3D manner










Computer-driven, Rapid Evaporative Imaging MS (REIMS) for tissue sections


23

Examining tissue (slices) by REIMS




Golf et al., Anal Chem 2015

Modes of data acquisition for REIMS







Individual Pixels: Pointing Mode




Golf et al., Anal Chem 2015

25









The IDH story of brain and other tumors

- IDH1 (isocitrate dehydrogenase) is mutated in position 132 in a GWAS study of patients with glioblastomas
- IDH1 catalyzes the conversion of isocitrate to alpha-ketoglutarate (α KG) which is a two-step reaction
- Mutant IDH1 catalyzes the first step to 2-hydroxyglutarate (2HG), but not the second one to α KG
- 2HG is considered to be an onco-metabolite
- Note that it has two stereoisomers
- What follows is a study from a group at Harvard performed in the Advanced Multimodality Image Guided Operating Suite at Brigham and Women's Hospital





Tumor xenograft imaging and 2HG

The ion at m/z 146.9 was subjected to MSMS to measure 2HG

IDH1 wt GBM (BT329)

IDH1 R132H GBM (BT116)


b


M21469


Santagata et al. PNAS 2014


Application to human glioblastoma m/z 147.2 1 mm 1 mm


Santagata et al. PNAS 2014


























Distribution of glucose/citrate ratio of some representative prostate tissue specimens showing significant elevation of the glucose/citrate ratio in cancer. The Top of each panel (A–O) shows the histopathological evaluation (H&E) of the corresponding tissue, where cancer areas have been demarcated by red, benign areas by black, stroma areas by green, and inflammation areas by blue.

43

Use of Raman spectroscopy Real-time imaging of metabolites in skin

• http://harvard.sunneyxielab.org/research/carstechniques.htm




Sunny Xie, PhD - ex-Harvard, now in Beijing

https://www.sunneyxielab.org/en_home/


Where to next?



MALDI-Imaging of a phospholipid Janusz Kabarowski/Kelly Walters



Multiple sampling single cells – Nemes group




CARS imaging of a cancer cell spectroscopic, real time Raman imaging

OR, two people with disparate abilities and insights will create something we've never heard of (yet)

45


What might it be?



Nanoprobe inserted into the wall of a cell recording changes in metabolism in real time sub nl sampling/analysis

Measuring O2 uptake using a Warburg apparatus - 10 ml incubations

YESTERDAY



The O₂ and pH probes of a Seahorse™ apparatus – 7 μl incubations

TODAY

TOMORROW?

Publications

- Santagata S, Eberlin LS, Norton I, Calligaris D, Feldman DR, Ide JL, Liu X, Wiley JS, Vestal ML, Ramkissoon SH, Orringer DA, Gill KK, Dunn IF, Dias-Santagata D, Ligon KL, Jolesz FA, Golby AJ, Cooks RG, Agar NY.
 Intraoperative mass spectrometry mapping of an onco-metabolite to guide brain tumor surgery.
 PNAS 2014;111(30):11121-6.
- Golf O, Strittmatter N, Karancsi T, Pringle SD, Speller AV, Mroz A, Kinross JM, Abbassi-Ghadi N, Jones EA, Takats Z. Rapid evaporative ionization mass spectrometry imaging platform for direct mapping from bulk tissue and bacterial growth media. Anal Chem. 2015 Mar 3;87(5):2527-34.
- Balog J, Kumar S, Alexander J, Golf O, Huang J, Wiggins T, Abbassi-Ghadi N, Enyedi A, Kacska S, Kinross J, Hanna GB, Nicholson JK, Takats Z. In vivo endoscopic tissue identification by rapid evaporative ionization mass spectrometry (REIMS). Angew Chem Int Ed Engl. 2015 Sep 14;54(38):11059-62.
- Banerjee S, Zarea RN, Tibshirani RJ, Kunder CA, Nolley R, Fan R, Brooks JD, Sonn GA. Diagnosis of prostate cancer by desorption electrospray ionization mass spectrometric imaging of small metabolites and lipids. PNAS early edition, March 2017
- St John ER, Balog J, McKenzie JS, Rossi M, Covington A, Muirhead L, Bodai Z, Rosini F, Speller AVM, Shousha S, Ramakrishnan R, Darzi A, Takats Z, Leff DR. Rapid evaporative ionisation mass spectrometry of electrosurgical vapours for the identification of breast pathology: towards an intelligent knife for breast cancer surgery. Breast Cancer Res. 2017 May 23;19(1):59.
- Phelps DL, Balog J, Gildea LF, Bodai Z, Savage A, El-Bahrawy MA, Speller AV, Rosini F, Kudo H, McKenzie JS, Brown R, Takáts Z, Ghaem-Maghami S. The surgical intelligent knife distinguishes normal, borderline and malignant gynaecological tissues using rapid evaporative ionisation mass spectrometry (REIMS). Br J Cancer. 2018 May;118(10):1349-1358.

47

More publications

- Alfaro CM, Pirro V, Keating MF, Hattab EM, Cooks RG, Cohen-Gadol AA. Intraoperative
 assessment of isocitrate dehydrogenase mutation status in human gliomas using
 desorption electrospray ionization-mass spectrometry. <u>J Neurosurg. 2019 Jan 4;132(1):180-187</u>.
- Pu F, Alfaro CM, Pirro V, Xie Z, Ouyang Z, Cooks RG. Rapid determination of isocitrate dehydrogenase mutation status of human gliomas by extraction nanoelectrospray using a miniature mass spectrometer. <u>Anal Bioanal Chem. 2019 Feb 2. doi: 10.1007/s00216-019-01632-5</u>.
- Hänel L, Kwiatkowski M, Heikaus L, Schlüter H. Mass spectrometry-based intraoperative tumor diagnostics. Future Sci OA. 2019 Mar 7;5(3):FSO373.
- Tzafetas M, Mitra A, Paraskevaidi M, Bodai Z, Kalliala I, Bowden S, Lathouras K, Rosini F, Szasz M, Savage A, Balog J, McKenzie J, Lyons D, Bennett P, MacIntyre D, Ghaem-Maghami S, Takats Z, Kyrgiou M. The intelligent knife (iKnife) and its intraoperative diagnostic advantage for the treatment of cervical disease. Proc Natl Acad Sci U S A. 2020 Mar 16.
- Kuzmin AN, Pliss A, Rzhevskii A, Lita A, Larion M. BCAbox Algorithm Expands Capabilities of Raman Microscope for Single Organelles Assessment. <u>Biosensors (Basel)</u>. 2018 Nov 10;8(4):106
- Ruiz-Rodado V, Lita A, Larion M. Advances in measuring cancer cell metabolism with subcellular resolution. <u>Nat Methods. 2022 Sep;19(9):1048-1063</u>.